

UCRL- 93603
PREPRINT

TAU ELECTRON ATOMS AT RHIC

Morton S. Weiss
University of California
Lawrence Livermore National Laboratory
Livermore, CA 94550

This paper was prepared for submittal to the
International Centre for Theoretical
Physics Topical Meeting
Trieste, Italy
September 30 - October 4, 1985

October 1985

The logo of the Lawrence Livermore National Laboratory is a large, stylized 'V' shape. The top horizontal bar of the 'V' is white with a stippled texture. The two slanted sides of the 'V' are dark grey with a horizontal line pattern. On the right slanted side, the words 'Lawrence Livermore National Laboratory' are written in a white, sans-serif font, oriented diagonally to follow the slope of the 'V'.

Lawrence
Livermore
National
Laboratory

This is a preprint of a paper intended for publication in a journal or proceedings. Since changes may be made before publication, this preprint is made available with the understanding that it will not be cited or reproduced without the permission of the author.

DISCLAIMER

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial products, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement recommendation, or favoring of the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

TAU ELECTRON ATOMS AT RHIC

Morton S. Weiss
Physics Department
Lawrence Livermore National Laboratory
Livermore, CA, 94550

September 12, 1985

Invited paper prepared for "Phase Space Approach To Nuclear Dynamics", ICTP, 30 September 4 - October, 1985

An amusement ancillary to the proposed quark-gluon plasma production hypothesized from a relativistic heavy ion collider (RHIC) may be a sufficient quantity of tau electrons to potentially admit the study of its exotic atoms. Given the wealth of nuclear phenomena derived from muonic atoms one might assume a tau atom would be even more forthcoming of information due to its lower orbits being entirely contained within the nucleus. It is the purpose of this brief note to discuss the production mechanism at a RHIC and to delineate some of the more obvious properties of the tau atom. As in the case of the mu, no doubt a host of more exotic phenomena derived from resonance "accidents" with nuclear transitions will also take place, but it would be presumptuous to discuss them at this time. Given the complete containment in nuclear matter of the tau lepton in its innermost atomic orbits, one would think the most immediately interesting, albeit potentially fruitless, experiment to be performed with such an exotic species would be the measurement of its lifetime.

These simple speculations are motivated by the work of Harvey

Gould¹, who has calculated background processes at a RHIC. Among the plethora of peripheral processes will be the production of lepton pairs by "the photon field" of one nucleus interacting with the coulomb field of the other. (In a collider one cannot use the comforting target and projectile nomenclature and I do not know of a euphonic alternative.) For the counterrotating beams of fully stripped Uranium at 100 GeV/A design that Gould considers, he estimates 10^8 electron pairs, $2 \cdot 10^3$ muon pairs and 1 tau pair would be produced per second. These are undoubtedly upper limits in that he ignores finite size effects which clearly will be most important for the tau, with its mass of 1784.2 MeV. A lower bound would reduce his estimate by two orders of magnitude but one can make arguments that that clearly would be a gross exaggeration. In fact the question of realistically incorporating finite size effects and inelasticity is not at all trivial and I am not aware of any such published treatment. Hence with the caveat that what follows may be an amusing exercise in futility unless the luminosity of $10^{27} \text{ cm}^{-2} \text{ s}^{-1}$ that Gould uses as a design figure were increased to compensate for the potential finite size decrement, we will proceed to calculate the spectra of a negatively charged tau electron captured by Pb.

However before doing so, some of the relative time scales involved should be discussed lest we waste time on the energy levels of a quantity whose unproducibility is exacerbated by its ephemerality. With a weak decay lifetime of $(3.4 \pm 0.5) \cdot 10^{-13}$ seconds the inherent instability of the tau- is only an order of

magnitude longer than the assumed cascade time of a μ^- of 10^{-14} seconds. However, the μ^- spends a great deal of this time losing energy by the auger process² at very distant orbits from the well shielded nucleus. The τ^- , at a RHIC, is instead created in the field of a bare nucleus which will both facilitate its capture and lead it into interesting x-ray producing orbits more quickly than a μ^- . The other process which might vitiate the existence of such exotic tau atoms is inverse beta decay. For a μ^- this is bounded by rates of 10^6 to 10^8 per second. Taking the latter, appropriate for a heavy nucleus with the μ^- half inside the nuclear material, and scaling by phase space and a factor of two for complete immersion we are still three orders of magnitude short of the natural lifetime. If we ignore decreasing phase space as we proceed up the baryon energy scale, this must be multiplied by the number of baryons the proton could be converted into with the caveat that the baryon energy must be less than 3 GeV. Fortunately, this still leaves considerable room for new particles before this rate approaches the weak decay lifetime. However, the weak decay lifetime will produce a natural lower bound for the line width of the τ^- x-rays of approximately 0.1 eV. It is improbable that this limitation will seriously inconvenience experimenters in the 1990s trying to measure, as we will see, MeV x-rays, in a system moving at nearly the velocity of light.

The Bohr radius of a tau atom of lead is approximately 0.1 fm which is picuant but irrelevant given a nucleus 12 or 14 fm in diameter. We solve the non-relativistic Schroedinger equation, with

for orientation purposes, a uniform charge distribution with sharp cutoff in radius at $1.1 \cdot A^{1/3}$, which takes the well known form:

$$-\frac{\hbar^2}{2m} \Delta^2 \phi + V(r) \phi = \epsilon \phi \quad \text{where}$$

$$V(r) = \frac{Ze^2}{2R_0} \left(\left(\frac{r}{R_0} \right)^2 - 3 \right) ; r < R_0 ; = \frac{Ze^2}{r} : r > R_0$$

For simplicity we have taken the tau mass to be twice a nucleon mass and ignore any spin or spin-orbit effects. From Table I we see the results for the first six orbits for S, P and D states. Displayed are the eigenvalues and radii at which the wave functions have their maximum. One notes that only the lowest states are fully contained in the interior of the nucleus and impervious to the surface. This also manifests itself in Table II where the energies of the non-node changing transitions are calculated. $\hbar \omega$ is 2.9 Mev for the sharp cutoff model, with these parameters, and we see an harmonic spectrum only for the first orbits. As nodeless hydrogen orbits scale approximately as n^2 where n is the sum of radial quantum number + angular momentum +1, we cannot expect and do not see any hydrogen like spectra by $n=6$. In fact, beyond $n=6$, we are very close to the onset of a continuum in x-ray energies, which will include the hydrogen quasi continuum limit but probably not permit its separation from those states affected by the finite nuclear size.

For comparison, but not for realism, we have also calculated the same spectra using a woods-saxon charge distribution with the same rms radius as the sharp cut off model but with two values of the

surface diffusion, the conventional 0.65 f and 0.55 f. These results are contained in Table I-Model II, and Table I-Model III, respectively. Table II includes the predicted x rays from all three models.

One sees that the spectrum is nearly harmonic for all three models only for the lowest orbits (1d-1p, 1p-1s) and that the models differ among themselves at the 100 kev level for these transitions. This is clearly the only region insensitive to the nuclear shape or surface and provides access to the true interior of a nucleus (whether or not this would represent nuclear matter is a subject of speculation). The other transitions for the models differ by similar quantities. Given the expected result that some of the spectra will be sensitive to the nuclear shape will no doubt lead to the same type of sophisticated analysis that the muon has seen, if data are ever available. However, the weighting will be very different. Prior to designing an experimental program one would of course engage in the more detailed analysis requisite from coupling atomic and nuclear levels³ which will shift many of the levels calculated here by at least several percent. It is our purpose only to point out the gross features of the spectrum.

It is clear that the spin-orbit splitting will be quite small due to the large mass of the tau, approximaely 10^{-8} of the potential energy which is no more than 30 Mev. Alternatively, for the sharp cut-off potential (Model I), the effect of the spin-orbit force is to add a small linear term to the potential energy for a given j, which displaces the zero of the potential slightly. Again,

one sees that the effect is presumably unobservable with present techniques. Finally, examining the energy levels of a tau atom inside or adjacent to the high magnetic field due to the accelerator may offer some possibility for zeeman effect observations but that is too speculative even for this fantasy.

The calculations offered here were for Pb even though the presumed RHIC design is to accelerate Uranium. Uranium has its own special cachet when it comes to forming an exotic atom because of its deformation and fissibility, which would obscure the naive calculations presented here. However, its muonic atoms offer some very amusing properties⁴ and no doubt refined calculations of its tau atoms will be forthcoming of similarly enriching phenomena.

It is a pleasure to thank Harvey Gould, Arthur Kerman, Ray Sheline and Neal Snyderman for respectfully, stimulating, critical, encouraging and helpful discussions and specially James Spencer for lending his copy of Schiff's Quantum Mechanics without which all numbers calculated would be modulo random powers of four pi. Work performed under the auspices of the U.S. Department of Energy by the Lawrence Livermore National Laboratory under contract number W-7405-ENG-48.

REFERENCES

1. H. Gould, LBL-18593, November, 1984, available through National Technical Information Service, Springfield, VA 22161.
2. J. Hufner et. al. in V. W. Hughes and C. S. Wu, Muon Physics, Vol I, Academic Press, N.Y (1977).

3. G. A. Rinker and J. Speth, Nuclear Physics A306, 397, (1978).
4. E. Teller and M.S. Weiss, "Is the Muon a Multipole Meter?",
UCRL-83616, 1979, published in Festschrift for Maurice Goldhaber.

TABLE CAPTIONS

Table I: e is the eigenvalue in Mev in descending order for $L=0, 1$ and 2. Space is the total radial distance over which numerical integraton was performed. Max Wf is the distance from the origin at which the wave function has its maximum. Delta is the difference between adjacent levels of the same L . The three models repesent different assumptions about the shape of the nuclear charge distribution. They are I: sharp cutoff, radius of 6.545 f; II: Woods-Saxon shape with radius of 5.754 f and diffusness of 0.65 f; III: as II with radius 5.989 f and diffusness of 0.55 f. All three models have the same rms radius and total charge of 82 protons. The tau mass has been taken to be twice the average nucleon mass and center of mass corrections have not been made.

Table II: Transition energies in Mev as indicated by model described in Table I. The near equivelance of $nP-nS$ and $nD-nP$, where n is the radial quantum number, only for $n=1$ is a measure off the limits of the harmonic spectrum independent of the nuclear surface.

FIGURE CAPTION

Fig. 1: Sketch of the tau- atomic orbits superimposed upon the sharp cut off nuclear model (Model I). Labels refer to Table I.




FIGURE 1

TABLE I - MODEL I

THESE ARE FROM SHARP CUTOFF MODEL

L	n	e	space	max wf	delta
0	1	-22.63	16	2.6	
					5.804
	2	-16.826	16	5.4	4.94
	3	-11.886	24	8	3.237
	4	8.649	24	11.3	2.143
	5	-6.506	40	15.2	1.4572
	6	-5.0488	50	20	
1	1	-19.693	24	3.84	5.583
	2	-14.11	30	6.6	4.087
	3	-10.023	36	9.36	2.627
	4	-7.396	36	12.96	1.741
	5	-5.655	36	17.28	1.205
	6	-4.45	50	22.5	
2	1	-16.781	20	4.6	5.098
	2	-11.683	24	7.44	3.246
	3	-8.437	30	11.1	2.106
	4	-6.331	30	15	1.418
	5	-4.913	40	19.6	0.9953
	6	-3.9177	60	25.2	0.7217
	7	-3.196	60	31.2	

TABLE I - MODEL II

FROM WOODS SAXON SHAPE
 $R_0 = 5.754$, $a = 0.65$

L	n	e	space	max wf	delta
0	1	23.4555	12	2.52	6.3374
	2	17.1181	24	5.28	5.0241
	3	-12.094	24	7.92	3.3294
	4	8.7646	36	11.16	2.1845
	5	-6.5601	50	15	1.4812
	6	-5.0989	50	19.5	
1	1	-20.1635	24	3.84	5.855
	2	-14.3085	24	6.72	4.1446
	3	-10.1639	36	10.08	2.6768
	4	-7.4871	36	13.68	1.774
	5	-5.7131	50	18	1.2217
	6	-4.4914	50	23	
2	1	-16.9743	20	4.8	5.1435
	2	-11.8308	30	8.1	3.3116
	3	-8.5192	30	11.4	2.1334
	4	-6.3858	40	15.6	1.4336
	5	-4.9522	50	20.5	1.0026
	6	-3.9496	60	25.8	

TABLE I - MODEL III

FROM WOODS SAXON SHAPE
 $R_0 = 5.989$, $a = 0.55$

L	n	e	space	max wf	delta
0	1	23.215	12	2.52	
					6.1862
	2	17.0288	24	5.28	
					4.9924
	3	-12.0364	24	7.92	
					3.3057
	4	8.7307	36	11.16	
1					2.1739
	5	-6.5568	50	15	
					1.4728
	6	-5.084	50	19.5	
	1	-20.0287	24	3.84	
					5.7793
2	2	-14.2494	24	6.72	
					4.1256
	3	-10.1236	36	10.08	
					2.6618
	4	-7.4618	36	13.68	
					1.7658
	5	-5.696	50	18	
3					1.2167
	6	-4.4793	50	23	
	1	-16.917	20	4.8	
					5.1264
	2	-11.7906	30	8.1	
					3.2959
4	3	-8.4947	30	11.4	
					2.125
	4	-6.3697	40	15.6	
					1.4291
	5	-4.9406	50	20.5	
					1.0017
	6	-3.9389	60	25.8	

TABLE II

MODEL

TRANSITION	I	II	III
1p-1s	2.9370	3.2920	3.1863
1d-1p	2.910	3.1892	3.1117
2p-2s	2.7160	2.8096	2.7794
2d-2p	2.4270	2.4777	2.4588
3p-3s	1.8630	1.9301	1.9128
3d-3p	1.5860	1.6447	1.6289
4p-4s	1.2530	1.2775	1.2689
4d-4p	1.0650	1.1013	1.0921
5p-5s	0.8510	0.8670	0.8608
5d-5p	0.7420	0.7609	0.7554
6s-6p	0.5988	0.6075	0.6047
6d-dp	0.5323	0.5418	0.5404